

Laparoscopic Colorectal Surgery

Rapid Review and Update



R Sim

Centre for Advanced

Laparoscopic Surgery, TTSH



Scope

- Diagnostic Laparoscopy
- Laparoscopic Staging for Cancer
- Laparoscopic Appendicectomy
- Laparoscopic Colorectal Resection
- Laparoscopic Rectopexy



Diagnostic Laparoscopy


- Acute abdominal pain
- Chronic abdominal/pelvic pain syndromes
- Adhesions and intermittent bowel obs.
- Ascites
- Hernias
- GI bleed of obscure origin
- Local anaesthesia possible



Laparoscopic Staging for Cancer

- Pancreatic, gastric, ovarian cancers
- Carcinomatosis, Peritoneal cytology
- Liver biopsy
- Ultrasonography
- Palliative bypass possible








Laparoscopic Appendicectomy

- Particularly useful in women
- Proven in 39 RCTs, >4000 patients
- Halved wound infections, less pain POD1, reduced hospital stay and time to RTW
- 3x postop intraabd. abscesses especially if perforated
- Able to deal with unexpected findings laparoscopically

Laparoscopic Colorectal Resection

- Cancer, complicated diverticular disease, inflammatory bowel disease, functional
- Multiple quadrants
- Retract small bowel
- Expose and dissect large planes
- Remove large, bacteria-laden organ
- Malignancy, port site mets
- Perform bowel anastomosis



Short-term Quality-of-Life Outcomes Following Laparoscopic-Assisted Colectomy vs Open Colectomy for Colon Cancer

A Randomized Trial

Jane C. Weeks, MD

Heidi Nelson, MD

Shari Gelber, MS

Daniel Sargent, PhD

Georgene Schroeder, MS

for the Clinical Outcomes of Surgical Therapy (COST) Study Group

ALTHOUGH LAPAROSCOPIC TECHNIQUES were first described in 1901,¹ only in the past few years have newer optics and instrumentation allowed for the safe application of laparoscopic resection procedures. The first report of a successful laparoscopic cholecystectomy in 1987 was followed by rapid widespread adoption of the procedure.²⁻⁶ In recent years, laparoscopic procedures for a number of other nonmalignant abdominal diseases, including appendicitis, inguinal hernia, gastroesophageal reflux disease, hiatal hernia, and nonmalignant uterine conditions, have become routine. The interest in laparoscopic approaches for these conditions has been driven by the theoretical benefits, including reduced postoperative pain, shortened length of stay, and earlier return to work, and perhaps by the technological imperative.^{5,6}

Improvements in both technology and surgeons' comfort and skill with laparoscopic techniques have led to an inter-

Context Laparoscopic-assisted colectomy (LAC) has emerged as the preferred minimally invasive surgical strategy for diseases of the colon. The safety and efficacy of LAC for colon cancer are unknown, and the nature and magnitude of any quality-of-life (QOL) benefit resulting from LAC for colon cancer is also unknown.

Objective To compare short-term QOL outcomes after LAC vs open colectomy for colon cancer.

Design, Setting, and Participants Multicenter, randomized controlled trial (Clinical Outcomes of Surgical Therapy [COST]). Between September 1994 and February 1999, 37 of 48 centers provided data for the QOL component of the trial for 449 consecutive patients with clinically resectable colon cancer.

Main Outcome Measures Scores on the Symptoms Distress Scale (SDS), Quality of Life Index, and a single-item global rating scale at 2 days, 2 weeks, and 2 months postoperative; duration of postoperative in-hospital analgesic use; and length of stay.

Results Of 449 patients, 428 provided QOL data. In an intention-to-treat analysis comparing SDS pain intensity, SDS summary, QOL Index summary, and global rating scale scores at each time point, the only statistically significant difference observed between groups was the global rating scale score for 2 weeks postsurgery. The mean (median) global rating scale scores for 2 weeks postsurgery were 76.9 (80) for LAC vs 74.4 (75) for open colectomy ($P=.009$). While in the hospital, patients assigned to LAC required fewer days of both parenteral analgesics compared with patients assigned to open colectomy (mean [median], 3.2 [3] vs 4.0 [4] days; $P<.001$) and oral analgesics (mean [median], 1.9 [1] vs 2.2 [2] days; $P=.03$).

Conclusion Only minimal short-term QOL benefits were found with LAC for colon cancer compared with standard open colectomy. Until ongoing trials establish that LAC is as effective as open colectomy in preventing recurrence and death from colon cancer, this procedure should not be offered to patients with colon cancer.

JAMA. 2002;287:321-328

www.jama.com

est in extending the indications for laparoscopic surgery to include curative resection of colon cancer. In laparoscopic-assisted colectomy (LAC), mobilization of the bowel is conducted laparoscopically and then the bowel is externalized for resection and anastomosis. Laparoscopic-assisted colectomy has emerged as the preferred minimally invasive strategy for colonic

Conclusion Only minimal short-term QOL benefits were found with LAC for colon cancer compared with standard open colectomy. Until ongoing trials establish that LAC is as effective as open colectomy in preventing recurrence and death from colon cancer, this procedure should not be offered to patients with colon cancer.

Author Affiliations: Department of Adult Oncology, Dana-Farber Cancer Institute, Boston, Mass (Dr Weeks and Ms Gelber); and Departments of Surgery (Dr Nelson) and Biostatistics (Dr Sargent and Ms Schroeder), Mayo Clinic, Rochester, Minn. Drs Weeks and Nelson contributed equally to this article as co-chairs of the Writing Committee.

Members of the COST Study Group are listed at the end of this article.
Corresponding Author: Jane C. Weeks, MD, Dana-Farber Cancer Institute, 44 Binney St, Boston, MA 02115 (e-mail: jane_weeks@dfci.harvard.edu).
Reprints: Heidi Nelson, MD, Mayo Clinic, 200 First St SW, Rochester, MN 55905 (e-mail: nelsonh@mayo.edu).

See also p 377 and Patient Page.

Laparoscopy-assisted colectomy versus open colectomy for treatment of non-metastatic colon cancer: a randomised trial

Antonio M Lacy, Juan C García-Valdecasas, Salvadora Delgado, Antoni Castells, Pilar Taurá, Josep M Piqué, Josep Visa

Summary

Background Although early reports on laparoscopy-assisted colectomy (LAC) in patients with colon cancer suggested that it reduces perioperative morbidity, its influence on long-term results is unknown. Our study aimed to compare efficacy of LAC and open colectomy (OC) for treatment of non-metastatic colon cancer in terms of tumour recurrence and survival.

Methods From November, 1993, to July, 1998, all patients with adenocarcinoma of the colon were assessed for entry in this randomised trial. Adjuvant therapy and postoperative follow-up were the same in both groups. The main endpoint was cancer-related survival. Data were analysed according to the intention-to-treat principle.

Findings 219 patients took part in the study (111 LAC group, 108 OC group). Patients in the LAC group recovered faster than those in the OC group, with shorter peristalsis-detection ($p=0.001$) and oral-intake times ($p=0.001$), and shorter hospital stays ($p=0.005$). Morbidity was lower in the LAC group ($p=0.001$), although LAC did not influence perioperative mortality. Probability of cancer-related survival was higher in the LAC group ($p=0.02$). The Cox model showed that LAC was independently associated with reduced risk of tumour relapse (hazard ratio 0.39, 95% CI 0.19–0.82), death from any cause (0.48, 0.23–1.01), and death from a cancer-related cause (0.38, 0.16–0.91) compared with OC. This superiority of LAC was due to differences in patients with stage III tumours ($p=0.04$, $p=0.02$, and $p=0.006$, respectively).

Interpretation LAC is more effective than OC for treatment of colon cancer in terms of morbidity, hospital stay, tumour recurrence, and cancer-related survival.

Lancet 2002; **359**: 2224–29

Departments of Surgery (A M Lacy MD, J C García-Valdecasas MD, S Delgado MD, J Visa MD), **Gastroenterology** (A Castells MD, J M Piqué MD), and **Anaesthesia** (P Taurá MD), **Institut de Malalties Digestives, Hospital Clínic, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), University of Barcelona, 08036 Barcelona, Spain**

Correspondence to: Dr Antonio M Lacy (e-mail: alacy@medicina.ub.es)

Introduction

Colorectal cancer is the second leading cause of cancer-related death in Western countries. Prognosis associated with this disease has improved due to early diagnosis and changes in medical therapy. Adjuvant chemotherapy in colon cancer, radiotherapy, and introduction of the total mesorectal excision technique in rectal cancer have increased survival, especially in patients with stage III tumours. Moreover, oxaliplatin and irinotecan have improved the prognosis associated with metastatic colorectal cancer.¹

Laparoscopic surgery has led to great progress in the treatment of many gastrointestinal diseases.² Early reports on laparoscopy-assisted colectomy (LAC) in patients with colon cancer suggest that it lowers surgical trauma, decreases perioperative complications, and leads to more rapid recovery.^{3–6} However, development of port-site metastases in some cases showed that this approach was questionable.^{7,8}

Few preliminary data that compare LAC with open colectomy (OC) in colon cancer have been reported. They suggest that LAC is associated with reduced perioperative morbidity and very low risk of wound metastasis.^{4,6,9,10} However, there are no studies that compare LAC and OC in terms of tumour recurrence and survival.

In this article we report the results of a randomised trial in patients with non-metastatic colon cancer. The aim of the trial was to assess whether there are differences in cancer-related survival between LAC and OC.

Methods

Patients

From November, 1993, to July, 1998, all patients admitted to our unit with adenocarcinoma of the colon, 15 cm above the anal verge, were assessed. Exclusion criteria were: cancer located at the transverse colon, distant metastasis, adjacent organ invasion, intestinal obstruction, past colonic surgery, and no consent to participate in the study.

Randomisation was done the day before surgery. Patients were stratified in two groups according to tumour location (right or left side, with respect to the splenic flexure), and subsequently assigned to LAC or OC by means of sealed opaque envelopes containing computer-generated random numbers. To prevent selection bias, random numbers were generated by an investigator (AC) who was not involved in enrolment of participants.




Due to the limited evidence about LAC at the beginning of the study, interim analyses that assessed early morbidity, tumour recurrence, and port-site metastasis were planned during the first period.^{9,10} The study was approved by the institutional ethics of research committee and oral consent was obtained from each patient.

Interpretation LAC is more effective than OC for treatment of colon cancer in terms of morbidity, hospital stay, tumour recurrence, and cancer-related survival.

This superiority of LAC was due to differences in patients with stage III tumours

Laparoscopic Rectopexy

- Ideal laparoscopic operation
- Functional surgery
- No extraction wound
- Analogous to fundoplication
- Trauma of access
- Trauma of surgery





Conclusion

We have to work harder
so that patients heal better